

Article ID:1005-3085(2010)06-1111-07

Existence of Periodic Solutions for Recurrent Cellular Neural Networks with Distributed Delays*

ZHANG Hong-li¹, CHEN Fang-qi^{1,2}

(1- College of Science, Nanjing University of Aeronautics and Astronautics, Nanjing 210016;

2- Tianjin Key Laboratory of Nonlinear Dynamics and Chaos Control, Tianjin 300072)

Abstract: By means of the Mawhin continuation theorem, the existence of periodic solutions for recurrent neural networks is studied. We assume that the behaved function is located in a strip region, and the activation function is in a region between two linear functions.

Keywords: recurrent cellular neural networks; periodic solution; coincidence degree

Classification: AMS(2000) 47H11; 46B50 **CLC number:** O177.92; O175.14 **Document code:** A

1 Introduction

Cellular neural networks (CNNs) are introduced by Chua and Yang. They have many applications in optimization, associative memories, and signal processing. Recently, they have been studied in the literature^[1-3]. Because neural networks usually have a spatial extent due to the presence of a multitude of parallel pathways with a variety of axon sizes and lengths, there will be a distribution of propagation delays^[4]. Recently, recurrent neural networks have been widely studied. The Hopfield neural networks and cellular neural networks^[5] are representative of them.

In the literature^[6], the authors proposed a class of recurrent neural networks as follows

$$x'_i(t) = -f_i(x_i(t)) + \sum_{j=1}^n a_{ij}(t)g_j(x_j(t)) + \sum_{j=1}^n b_{ij}(t)g_j(x_j(t - \tau_{ij}(t))) + I_i(t), \quad t > 0,$$

$$x_i(t) = \phi_i(t) \neq 0, \quad t \in [-\tau, 0], \quad i = 1, 2, \dots, n,$$

where n denotes the number of units in neural networks, $x_i(t)$ is the state function of the i th unit, $a_{ij}(t)$ and $b_{ij}(t)$ denote the strength of connectivity between the cell i and j at time t , and they are both continuous ω -periodic functions, $f(x) = (f_1(x_1), f_2(x_2), \dots, f_n(x_n))^T$ is an appropriately behaved function, $g(x) = (g_1(x_1), \dots, g_n(x_n))^T$ is a nonlinear vector-valued activation function, $I(t) = (I_1(t), I_2(t), \dots, I_n(t))^T$ is a continuous ω -periodic vector-valued input function, and $0 \leq \tau_{ij}(t) \leq \tau$ is a time delay required in processing and transmitting a signal from the j th cell to the i th cell at time t .

Received: 16 Oct 2008.

Biography: Zhang Hongli (Born in 1982), Female, Master.

Accepted: 25 May 2009.

Research field: dynamical system.

*Foundation item: The National Natural Science Foundation of China (10972099); the Natural Science Foundation of Tianjin (09JCZDJC26800).

The aim of this paper is to investigate the existence of periodic solution for the networks

$$x'_i(t) = -f_i(x_i(t)) + \sum_{j=1}^n a_{ij}(t)g_j(x_j(t)) + \sum_{j=1}^n b_{ij}(t)g_j(x_j(t - \tau_{ij}(t))) \\ + \sum_{j=1}^n c_{ij}(t) \int_0^\infty k_{ij}(s)h_j(x_j(t-s))ds + I_i(t), \quad t \in [0, +\infty), \quad i = 1, 2, \dots, n. \quad (1)$$

In (1), f_i and g_i are continuous functions, c_{ij} is a continuous ω -periodic connection weight function, h_j is an activation function of signal transmission, and k_{ij} is a delay kernel function. The initial conditions associated with (1) are $x_i(s) = \phi_i(s) \neq 0$, $s \in (-\infty, 0]$, where $\phi_i \in C((-\infty, 0], \mathbf{R})$, $i = 1, 2, \dots, n$.

Throughout this paper, we assume that:

(H₁) There exist positive constants $\alpha'_i, \beta'_i, \alpha_i, \beta_i$ and $\beta'_i < \beta_i$ ($i = 1, 2, \dots, n$) such that

$$f_i(0) = 0, \quad \alpha'_i|u| - \beta'_i \leq f_i(u) \leq \alpha_i|u| + \beta_i, \quad u \in \mathbf{R}.$$

(H₂) There exist positive constants σ_j, θ_j such that

$$|g_j(u)| \leq \sigma_j|u| + \theta_j, \quad u \in \mathbf{R}, \quad j = 1, 2, \dots, n.$$

(H₃) There exist positive constants $Q_j > 0$ such that

$$|h_j(u)| \leq Q_j, \quad u \in \mathbf{R}, \quad j = 1, 2, \dots, n.$$

(H₄) $\tau_{ij}(t)$ ($i, j = 1, 2, \dots, n$) are continuously differentiable ω -periodic functions defined on $[0, +\infty)$ and

$$\inf_{t \in [0, +\infty)} (1 - \tau'_{ij}(t)) > 0.$$

(H₅) The delay kernels $k_{ij}(s) : [0, \infty) \rightarrow [0, \infty)$ are continuous, and

$$\int_0^\infty k_{ij}(s)ds = k_{ij} > 0, \quad i, j = 1, 2, \dots, n.$$

For the sake of convenience, we introduce the following notations

$$a_{ij}^+ = \max_{0 \leq t \leq \omega} |a_{ij}(t)|, \quad b_{ij}^+ = \max_{0 \leq t \leq \omega} |b_{ij}(t)|, \quad c_{ij}^+ = \max_{0 \leq t \leq \omega} |c_{ij}(t)|, \quad I_i^+ = \max_{0 \leq t \leq \omega} |I_i(t)|,$$

$$|\tilde{I}_i| = \frac{1}{\omega} \int_0^\omega |I_i(t)|dt, \quad m_{ij} = \left(\max_{0 \leq t \leq \omega} \frac{1}{1 - \tau'_{ij}(t)} \right)^{\frac{1}{2}}, \quad i, j = 1, 2, \dots, n.$$

2 Preliminaries

Firstly, we introduce the Mawhin continuation theorem^[7] as follows:

Lemma 2.1 Let X and Y be two Banach spaces, $L : \text{Dom } L \subset X \rightarrow Y$ be a Fredholm operator with index zero, $\Omega \subset X$ be an open bounded set, and $N : X \rightarrow Y$ be a continuous operator which is L -compact on $\bar{\Omega}$. Assume that

- (a) For each $\lambda \in (0, 1)$, $x \in \partial\Omega \cap \text{Dom } L$, $Lx \neq \lambda Nx$;
- (b) For each $x \in \partial\Omega \cap \text{Ker } L$, $QNx \neq 0$;
- (c) $\deg\{JQN, \Omega \cap \text{Ker } L, 0\} \neq 0$.

Then $Lx = Nx$ has at least one solution in $\bar{\Omega} \cap \text{Dom } L$.

3 Existence of periodic solutions

Theorem 3.1 Assume $(H_1)-(H_5)$ hold, and

$$\alpha'_i(1 - \alpha_i\omega) - \sigma_i \sum_{j=1}^n (1 + \alpha'_j\omega)(a_{ji}^+ + b_{ji}^+ m_{ji}) > 0,$$

$$\alpha'_i - \sum_{j=1}^n (a_{ji}^+ + b_{ji}^+) \sigma_i > 0, \quad i = 1, 2, \dots, n,$$

then the network (1) has at least one ω -periodic solution.

Proof Let

$$X = Z = \{x(t) \in C(\mathbf{R}, \mathbf{R}^n) : x(t) = x(t + \omega), x = (x_1, x_2, \dots, x_n)^T\}.$$

Then X and Z are both Banach spaces with the norm

$$\|x\|_2 = \left(\int_0^\omega |x(t)|^2 dt \right)^{\frac{1}{2}}.$$

Let $L : \text{Dom } L \subset X \rightarrow Z$, $Lx = x'$, $x \in \text{Dom } L \subset X$, where $\text{Dom } L = C^1(\mathbf{R}, \mathbf{R}^n) \cap X$ and $N : X \rightarrow Z$,

$$Nx = \left(-f_i(x_i(t)) + \sum_{j=1}^n a_{ij}(t)g_j(x_j(t)) + \sum_{j=1}^n b_{ij}(t)g_j(x_j(t - \tau_{ij}(t))) \right. \\ \left. + \sum_{j=1}^n c_{ij}(t) \int_0^\infty k_{ij}(s)h_j(x_j(t-s))ds + I_i(t) \right)_{n \times 1}.$$

Obviously, $\text{Im } L$ is closed in Z and L is a Fredholm mapping of index zero. Define two projectors P and Q as

$$Px = \frac{1}{\omega} \int_0^\omega x(t) dt, \quad x \in X, \quad Qz = \frac{1}{\omega} \int_0^\omega z(t) dt, \quad z \in Z.$$

So, N is L -compact on $\bar{\Omega}$ for any open bounded set $\Omega \subset X$.

Corresponding to the operator equation $Lx = \lambda Nx$, $\lambda \in (0, 1)$, we have

$$\frac{dx_i(t)}{dt} = \lambda \left[-f_i(x_i(t)) + \sum_{j=1}^n a_{ij}(t)g_j(x_j(t)) + \sum_{j=1}^n b_{ij}(t)g_j(x_j(t - \tau_{ij}(t))) \right. \\ \left. + \sum_{j=1}^n c_{ij}(t) \int_0^\infty k_{ij}(s)h_j(x_j(t-s))ds + I_i(t) \right], \quad i = 1, 2, \dots, n. \quad (2)$$

Suppose that $x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T \in X$ is a solution to (2) for some $\lambda \in (0, 1)$.

Multiplying both sides of (2) by x'_i and integrating over $[0, \omega]$, we obtain

$$\begin{aligned} \int_0^\omega |x'_i|^2 dt &\leq \alpha_i \|x'_i\|_2 \|x_i\|_2 + \beta_i \sqrt{\omega} \|x'_i\|_2 + \sum_{j=1}^n a_{ij}^+ \sigma_j \|x'_i\|_2 \|x_j\|_2 \\ &\quad + \sum_{j=1}^n a_{ij}^+ \theta_j \sqrt{\omega} \|x'_i\|_2 + I_i^+ \sqrt{\omega} \|x'_i\|_2 + \sum_{j=1}^n b_{ij}^+ \sigma_j m_{ij} \|x'_i\|_2 \|x_j\|_2 \\ &\quad + \sum_{j=1}^n b_{ij}^+ \theta_j \sqrt{\omega} \|x'_i\|_2 + \sum_{j=1}^n c_{ij}^+ k_{ij} Q_j \sqrt{\omega} \|x'_i\|_2. \end{aligned} \quad (3)$$

From (3), it follows that

$$\begin{aligned} \|x'_i\|_2 &\leq \alpha_i \|x_i\|_2 + \sum_{j=1}^n (a_{ij}^+ \sigma_j + b_{ij}^+ \sigma_j m_{ij}) \|x_j\|_2 + \left(\beta_i \sqrt{\omega} + \sum_{j=1}^n a_{ij}^+ \theta_j \sqrt{\omega} \right. \\ &\quad \left. + \sum_{j=1}^n b_{ij}^+ \theta_j \sqrt{\omega} + \sum_{j=1}^n c_{ij}^+ k_{ij} Q_j \sqrt{\omega} + I_i^+ \sqrt{\omega} \right), \quad i = 1, 2, \dots, n. \end{aligned}$$

Integrating both sides of (1) over $[0, \omega]$, we obtain

$$\begin{aligned} \int_0^\omega f_i(x_i(t)) dt &= \sum_{j=1}^n \int_0^\omega a_{ij}(t) g_j(x_j(t)) dt + \sum_{j=1}^n \int_0^\omega b_{ij}(t) g_j(x_j(t - \tau_{ij}(t))) dt \\ &\quad + \int_0^\omega I_i(t) dt + \sum_{j=1}^n \int_0^\omega c_{ij}(t) \int_0^\infty k_{ij}(s) h_j(x_j(t-s)) ds dt. \end{aligned}$$

Then, there exists some $\xi \in (0, \omega)$ such that

$$\begin{aligned} f_i(x_i(\xi)) &\leq \frac{1}{\omega} \sum_{j=1}^n a_{ij}^+ \int_0^\omega (\sigma_j |x_j(t)| + \theta_j) dt \\ &\quad + \frac{1}{\omega} \sum_{j=1}^n b_{ij}^+ \int_0^\omega (\sigma_j |x_j(t - \tau_{ij}(t))| + \theta_j) dt + \sum_{j=1}^n c_{ij}^+ k_{ij} Q_j + I_i^+. \end{aligned} \quad (4)$$

Applying (H_1) to inequality (4) and by the Hölder inequality, we have

$$\begin{aligned} \alpha'_i |x_i(\xi)| &\leq \frac{1}{\sqrt{\omega}} \sum_{j=1}^n (a_{ij}^+ + b_{ij}^+ m_{ij}) \sigma_j \|x_j\|_2 \\ &\quad + \sum_{j=1}^n a_{ij}^+ \theta_j + \sum_{j=1}^n b_{ij}^+ \theta_j + \sum_{j=1}^n c_{ij}^+ k_{ij} Q_j + I_i^+ + \beta'_i. \end{aligned}$$

For any $t \in [0, \omega]$, it follows that

$$|x_i(t)| \leq |x_i(\xi)| + \int_0^\omega |x'_i(t)| dt \leq |x_i(\xi)| + \sqrt{\omega} \|x'_i\|_2, \quad i = 1, 2, \dots, n.$$

Consequently

$$\alpha'_i \|x_i\|_2 = \alpha'_i \left(\int_0^\omega |x_i(t)|^2 dt \right)^{\frac{1}{2}} \leq \alpha'_i \sqrt{\omega} \max_{0 \leq t \leq \omega} |x_i(t)| \leq \alpha'_i \sqrt{\omega} |x_i(\xi)| + \alpha'_i \omega \|x'_i\|_2.$$

Then

$$\alpha'_i (1 - \alpha_i \omega) \|x_i\|_2 \leq (1 + \alpha'_i \omega) \sum_{j=1}^n (a_{ij}^+ + b_{ij}^+ m_{ij}) \sigma_j \|x_j\|_2 + R_i^*, \quad (5)$$

where

$$\begin{aligned} R_i^* \equiv & \sqrt{\omega} \left(\sum_{j=1}^n a_{ij}^+ \theta_j + \sum_{j=1}^n b_{ij}^+ \theta_j + \sum_{j=1}^n c_{ij}^+ k_{ij} Q_j + I_i^+ + \beta'_i \right) \\ & + \alpha'_i \omega \left(\beta_i \sqrt{\omega} + \sum_{j=1}^n a_{ij}^+ \theta_j \sqrt{\omega} + \sum_{j=1}^n b_{ij}^+ \theta_j \sqrt{\omega} + \sum_{j=1}^n c_{ij}^+ k_{ij} Q_j \sqrt{\omega} + I_i^+ \sqrt{\omega} \right). \end{aligned}$$

From (5), it follows that

$$\sum_{i=1}^n \alpha'_i (1 - \alpha_i \omega) \|x_i\|_2 \leq \sum_{i=1}^n \left((1 + \alpha'_i \omega) \sum_{j=1}^n (a_{ij}^+ + b_{ij}^+ m_{ij}) \sigma_j \|x_j\|_2 + R_i^* \right).$$

Consequently

$$\sum_{i=1}^n \|x_i\|_2 \leq H^*,$$

where

$$H^* \equiv \frac{\sum_{i=1}^n R_i^*}{\min_{1 \leq i \leq n} [\alpha'_i (1 - \alpha_i \omega) - \sigma_i \sum_{j=1}^n (1 + \alpha'_j \omega) (a_{ji}^+ + b_{ji}^+ m_{ji})]}.$$

Clearly, H^* is independent of λ . Denote $M^* = H^* + F$, where $F > 0$ is sufficiently large so that

$$\min_{1 \leq i \leq n} \left[\alpha'_i - \sum_{j=1}^n (a_{ji}^+ + b_{ji}^+) \sigma_i \right] * \frac{M^*}{\sqrt{\omega}} - \sum_{i=1}^n \sum_{j=1}^n [(a_{ij}^+ + b_{ij}^+) \theta_j + c_{ij}^+ Q_j k_{ij}] - \sum_{i=1}^n (\beta'_i + |\tilde{I}_i|) > 0.$$

Now we define

$$\Omega = \left\{ x = (x_1(t), x_2(t), \dots, x_n(t))^T \in X \mid \sum_{i=1}^n \|x_i\|_2 < M^* \right\}.$$

Then Ω satisfies condition (a) in Lemma 2.1. Let $(x_1, x_2, \dots, x_n)^T \in \partial\Omega \cap \text{Ker } L$, $(x_1, x_2, \dots, x_n)^T$ is a constant vector in \mathbf{R}^n with

$$\|(x_1, x_2, \dots, x_n)\|_0 \equiv |x_1| + |x_2| + \dots + |x_n| = \frac{M^*}{\sqrt{\omega}}.$$

Then

$$\begin{aligned}
 \|QN(x_1, x_2, \dots, x_n)\|_0 &\geq \sum_{i=1}^n |f_i(x_i)| - \sum_{i=1}^n \left| \sum_{j=1}^n (a_{ij}^+ g_j(x_j) + b_{ij}^+ g_j(x_j)) \right| \\
 &\quad - \frac{1}{\omega} \sum_{i=1}^n \int_0^\omega |I_i(t)| dt - \sum_{i=1}^n \sum_{j=1}^n |c_{ij}^+ h_j(x_j) k_{ij}| \\
 &\geq \min_{1 \leq i \leq n} \left[\alpha'_i - \sum_{j=1}^n (a_{ji}^+ + b_{ji}^+) \sigma_i \right] \sum_{i=1}^n |x_i| \\
 &\quad - \sum_{i=1}^n \sum_{j=1}^n [(a_{ij}^+ + b_{ij}^+) \theta_j + c_{ij}^+ Q_j k_{ij}] - \sum_{i=1}^n (\beta'_i + |\tilde{I}_i|) > 0.
 \end{aligned}$$

This implies condition (b) of Lemma 2.1.

Define $\Psi : \text{Ker } L \times [0, 1] \rightarrow X$ by

$$\begin{aligned}
 \Psi(x_1, x_2, \dots, x_n, \mu) &= -\mu \left(\frac{1}{\omega} \int_0^\omega f_1(x_1) dt, \frac{1}{\omega} \int_0^\omega f_2(x_2) dt, \dots, \frac{1}{\omega} \int_0^\omega f_n(x_n) dt \right)^T \\
 &\quad + (1 - \mu) QN(x_1, x_2, \dots, x_n)^T, \\
 \|\Psi(x_1, x_2, \dots, x_n, \mu)\|_0 &\geq \sum_{i=1}^n |f_i(x_i)| - \sum_{i=1}^n \left[\sum_{j=1}^n (a_{ij}^+ + b_{ij}^+) (\sigma_j |x_j| + \theta_j) + \sum_{j=1}^n c_{ij}^+ Q_j k_{ij} + |\tilde{I}_i| \right] \\
 &\geq \min_{1 \leq i \leq n} \left[\alpha'_i - \sum_{j=1}^n (a_{ji}^+ + b_{ji}^+) \sigma_i \right] \sum_{i=1}^n |x_i| \\
 &\quad - \sum_{i=1}^n \sum_{j=1}^n [(a_{ij}^+ + b_{ij}^+) \theta_j + c_{ij}^+ Q_j k_{ij}] - \sum_{i=1}^n (\beta'_i + |\tilde{I}_i|) > 0.
 \end{aligned}$$

Hence, using the homotopy invariance theorem, we obtain

$$\deg(QN(x_1, x_2, \dots, x_n)^T, \Omega \cap \text{Ker } L, (0, 0, \dots, 0)^T) \neq 0.$$

By Lemma 2.1, (1) has at least one ω -periodic solution.

4 Conclusions

By using the Mawhin continuation theorem, derived in this paper are some new sufficient conditions ensuring the existence of periodic solutions for recurrent cellular neural networks with distributed delays.

References:

- [1] 刘艳青, 唐万生. 带有周期系数和时滞的细胞神经网络模型的周期解存在性和全局指数稳定性[J]. 工程数学学报, 2007, 24(6): 995-1006
- Liu Y Q, Tang W S. Existence and global exponential stability of periodic solution of cellular neural networks with periodic coefficients and delays[J]. Chinese Journal of Engineering Mathematics, 2007, 24(6): 995-1006

- [2] 邱亚林. 具有时滞的细胞神经网络吸引集与周期解存在性[J]. 工程数学学报, 2008, 25(2): 211-218
Qiu Y L. Existence of attracting sets and periodic solutions for cellular neural networks with delays[J]. Chinese Journal of Engineering Mathematics, 2008, 25(2): 211-218
- [3] 蒋秋浩, 曹进德. 具有变时滞的神经网络的全局指数稳定性[J]. 工程数学学报, 2006, 23(2): 373-376
Jiang Q H, Cao J D. Delay-dependent global exponential stability analysis of delayed neural networks with time-varying delays[J]. Chinese Journal of Engineering Mathematics, 2006, 23(2): 373-376
- [4] Li K, Zhang X, Li Z. Global exponential stability of impulsive cellular neural networks with time-varying and distributed delay[J]. Chaos, Solitons and Fractals, 2009, 41(3): 1427-1434
- [5] Chua L O, Yang L. Cellular neural networks: theory[J]. IEEE Transactions on Circuits Systems, 1988, 35(10): 1257-1272
- [6] Sun C, Feng C. Exponential stability and periodicity of delayed neural networks[J]. Mathematics and Computers in Simulation, 2004, 66(6): 469-478
- [7] 郭大均, 孙经先, 刘兆理. 非线性常微分方程泛函方法[M]. 济南: 山东科学技术出版社, 2006
Guo D J, Sun J X, Liu Z L. Functional Methods in Nonlinear Ordinary Differential Equations[M]. Jinan: Shandong Science and Technology Press, 2006

具分布时滞的周期运动细胞神经网络周期解的存在性

张红丽¹, 陈芳启^{1,2}

(1- 南京航空航天大学理学院, 南京 210016;

2- 天津市非线性动力与混沌控制重点实验室, 天津 300072)

摘 要: 运用 Mawhin 连续性定理研究具分布时滞的周期运动细胞神经网络周期解的存在性, 假设行为函数位于一带型区域内, 激活函数位于两线性函数所夹的区域内。

关键词: 周期运动细胞神经网络; 周期解; 迭合度